


Assessment Schedule – 2005**Scholarship Mathematics with Calculus: (93202)****Evidence Statement**

As shown in the schedule below, either a seven-point marking scale (0–6), or a nine-point marking scale (0–8), was used to assess the questions.

Question	Evidence	Code	Judgement
ONE (a)	$z^5 - 1 = 0$ $\alpha^5 - 1 = 0$ $(\alpha - 1)(\alpha^4 + \alpha^3 + \alpha^2 + \alpha + 1) = 0$ but α is complex so $\alpha^4 + \alpha^3 + \alpha^2 + \alpha + 1 = 0$ $\alpha^4 + \alpha^3 + \alpha^2 + \alpha = -1$. Sum of roots is $\alpha^4 + \alpha^3 + \alpha^2 + \alpha = -1$ from above.	6	No simplification: –1 mark. No ‘hence’: –1 mark. no sum of roots: max. 4 marks.
	Product of roots is $(\alpha + \alpha^4)(\alpha^2 + \alpha^3) = \alpha^3 + \alpha^4 + \alpha^6 + \alpha^7$ But $\alpha^5 = 1$ $\alpha^3 + \alpha^4 + \alpha^6 + \alpha^7 = \alpha^3 + \alpha^4 + \alpha + \alpha^2 = -1$ hence the equation is: $z^2 + z - 1 = 0$.	2	
ONE (b)(i)	 Either As shown following, BE is perpendicular to the x -axis (by symmetry, congruent triangles, AXB and AXE, SAS).	6	

Question	Evidence	Code	Judgement
ONE (b)(i) contd	So $BE = 2b$, but $BE = CE$ (triangles ABE and DEC congruent SAS) So $CE = 2b$. Or $\angle BOA = \frac{2\pi}{5}$ (72°) and $OB = 2$, hence $b = 2\sin \frac{2\pi}{5}$. Triangle COE is isosceles ($CO = EO = 2$). So $CY = EY = 2\sin \frac{2\pi}{5} = b$ and so $CE = 2b$. Alternative for this last step is the cosine rule: $CE^2 = 2^2 + 2^2 - 2 \cdot 2 \cdot 2 \cos \frac{4\pi}{5}$ $CE^2 = 8 \left(1 - \cos \frac{4\pi}{5} \right)$ $= 8 \left(1 - (1 - 2 \sin^2 \frac{2\pi}{5}) \right) = 8 \left(2 \sin^2 \frac{2\pi}{5} \right) = 16 \sin^2 \frac{2\pi}{5}$ and $CE = 4 \sin \frac{2\pi}{5} = 2b$.		Accept degrees.
			Answer must be in terms of b . Accept decimals if = $2b$. If answer from decimals is only inferred: –1 mark.

Question	Evidence	Code	Judgement
ONE (b)(ii) contd	$= \sqrt{2\frac{1}{2}} + \sqrt{2}\sqrt{1\frac{1}{4}} = \sqrt{2\frac{1}{2}} + \sqrt{2\frac{1}{2}} = 2\sqrt{\frac{5}{2}} = \sqrt{10}.$ Or: Geometrically, let A' be the reflection of A in the line $y = x$ that W lies on. So $A' = (0,2)$. Then $AW = A'W$ and the minimum value of $A'W + FW$ is when A'F is a straight line (ie for the point W_2 as shown above). Here $A'W_2 + FW_2 = A'F$ and by Pythagoras' theorem $A'F = \sqrt{1^2 + 3^2} = \sqrt{10}, \text{ the minimum value.}$ Or: Gradient of AF = gradient of the line $y = x$ that W lies on. So they are parallel. Hence the minimum distance is when WAF is an isosceles triangle, and the line from W is perpendicular to AF. Hence $AW + FW = 2AW = 2FW$ Equation of AF is $y = x - 2$ and perpendicular distance between the lines $= 2 \sin 45^\circ = \sqrt{2}$ $AF^2 = 1^2 + (-1)^2 = 2 \text{ so}$ $AW^2 = \left(\sqrt{2}\right)^2 + \left(\frac{\sqrt{2}}{2}\right)^2$ and minimum $AW + FW = 2AW = 2\sqrt{\frac{5}{2}} = \sqrt{10}.$		Accept 3.162 or decimal equivalent.

TWO (a)	The equation of the circle $(x - a)^2 + (y - b)^2 = r^2$ The point $A(x, y) \rightarrow B(x', y')$ A, B correct $x' = x \qquad x = x'$ $y' = hy \qquad y = \frac{y'}{h}$ substituting gives $(x' - a)^2 + \left(\frac{y'}{h} - b\right)^2 = r^2$ $\frac{(x' - a)^2}{r^2} + \frac{(y' - hb)^2}{h^2 r^2} = 1$ $h = \frac{1}{2}$ equation of ellipse is $\frac{(x - a)^2}{r^2} + \frac{\left(y - \frac{1}{2}b\right)^2}{\frac{r^2}{4}} = 1$ Method 1 $(x - a)^2 + (2y - b)^2 = r^2$ on the y-axis $x = 0$ $a^2 + (2y - b)^2 = r^2$ $2y - b = \pm \sqrt{r^2 - a^2}$ $y = \frac{b \pm \sqrt{r^2 - a^2}}{2}$ The circle cuts the y-axis when $a^2 + (y - b)^2 = r^2$ $y - b = \pm \sqrt{r^2 - a^2}$ $y = b \pm \sqrt{r^2 - a^2}$ For the ellipse and the circle to intersect on the y-axis: $\frac{b + \sqrt{r^2 - a^2}}{2} = b - \sqrt{r^2 - a^2}$ $b + \sqrt{r^2 - a^2} = 2b - 2\sqrt{r^2 - a^2}$ $b = 3\sqrt{r^2 - a^2}$ $b^2 = 9(r^2 - a^2)$	6 4
----------------------------------	--	---------------------------------

TWO (a) contd	Method 2 Solve equations simultaneously: $(x-a)^2 + (2y-b)^2 = (x-a)^2 + (y-b)^2$ $2y-b = \pm(y-b)$ $2y-b = +y-b \quad \text{or} \quad 2y-b = -(y-b)$ $y = 0 \qquad y-b = -y+b$ or $y = \frac{2b}{3}$ To meet on the y-axis $x = 0$, so for the circle using $y = \frac{2b}{3}$ $(x-a)^2 + (y-b)^2 = r^2$ $a^2 + \left(\frac{2b}{3} - b\right)^2 = r^2$ $\left(\frac{b}{3}\right)^2 = r^2 - a^2$ $b^2 = 9(r^2 - a^2)$ Method 3 If the y co-ordinate of one point where the ellipse and the circle intersect on the y-axis is y_1 then the other is $0.5y_1$ For the circle, $(x-a)^2 + (y-b)^2 = r^2$, when $x = 0$, $y^2 - 2by + b^2 + a^2 - r^2 = 0$ but the sum of these roots is $1.5y_1$ and the product $0.5y_1^2$ so $\frac{3}{2}y_1 = 2b$ $\frac{1}{2}y_1^2 = b^2 + a^2 - r^2$ and so $\frac{1}{2}\left(\frac{16}{9}b^2\right) = b^2 + a^2 - r^2$, $\frac{1}{9}b^2 = r^2 - a^2$, $b^2 = 9(r^2 - a^2)$.	4	
TWO (b)	Circle: $x^2 + y^2 = r^2$ $y^2 = r^2 - x^2$ $y = \pm\sqrt{r^2 - x^2}$ Since we require the top half of the circle, $y = +\sqrt{r^2 - x^2}$	6	

TWO
(b)
contd

Ellipse:

$$x^2 + 16(y - r)^2 = r^2$$

Method 1

$$x^2 + y^2 = r^2$$

∴ Meet where (subtracting)

$$16(y - r)^2 - y^2 = 0$$

$$4(y - r) = \pm y$$

$$5y = 4r \text{ or } 3y = 4r$$

$$y = \frac{4r}{5} \text{ or } \frac{4r}{3}$$

but $y < r$

$$\therefore y = \frac{4r}{5}$$

$$\therefore x^2 = r^2 - \frac{16r^2}{25} = \frac{9r^2}{25}$$

$$x = \pm \frac{3r}{5}$$

2

Method 2

$$(y - r)^2 = \frac{r^2 - x^2}{16}$$

$$y - r = \pm \sqrt{\frac{r^2 - x^2}{16}}$$

$$y = r \pm \frac{\sqrt{r^2 - x^2}}{4}$$

For $y < r$

$$y = r - \frac{\sqrt{r^2 - x^2}}{4}$$

Solving for points of intersection:

TWO (b) Contd	$r - \frac{\sqrt{r^2 - x^2}}{4} = \sqrt{r^2 - x^2}$ $\sqrt{r^2 - x^2} \left(1 + \frac{1}{4}\right) = r$ $\sqrt{r^2 - x^2} = \frac{4r}{5}$ $r^2 - x^2 = \frac{16r^2}{25}$ $x^2 = \left(1 - \frac{16}{25}\right)r^2$ $= \frac{9r^2}{25}$ $x = \pm \frac{3r}{5}$	2	Accept decimal limit.
	$\text{Area} = 2 \int_0^{\frac{3r}{5}} \sqrt{r^2 - x^2} - \left(r - \frac{\sqrt{r^2 - x^2}}{4}\right) dx$ $= 2 \int_0^{\frac{3r}{5}} \frac{5}{4} \sqrt{r^2 - x^2} - r dx$ $= 2 \int_0^{\frac{3r}{5}} \frac{5}{4} \sqrt{r^2 - x^2} dx - 2 \int_0^{\frac{3r}{5}} r dx$ using the substitution: $x = r \sin u \quad x = 0 \quad u = 0$ $dx = r \cos u du \quad x = \frac{3r}{5} \quad u = \sin^{-1}\left(\frac{3}{5}\right) = 0.6435$ $2 \int_0^{\frac{3r}{5}} \frac{5}{4} \sqrt{r^2 - x^2} dx = 2 \int_0^{\sin^{-1}\left(\frac{3}{5}\right)} \frac{5}{4} \sqrt{(r^2 - r^2 \sin^2 u)} r \cos u du$		
	$= 2 \int_0^{\sin^{-1}\left(\frac{3}{5}\right)} \frac{5}{4} r \cos u r \cos u du$ $= 2 \times \frac{5}{4} r^2 \int_0^{\sin^{-1}\left(\frac{3}{5}\right)} \cos^2 u du$ $= \frac{5}{4} r^2 \int_0^{\sin^{-1}\left(\frac{3}{5}\right)} (\cos 2u + 1) du$	4	

	$= \frac{5}{4}r^2 \left[\frac{1}{2} \sin 2u + u \right]_0^{\sin^{-1}\left(\frac{3}{5}\right)}$ $= \frac{5}{4}r^2 \left[\sin u \cos u + u \right]_0^{\sin^{-1}\left(\frac{3}{5}\right)}$ $= \frac{5}{4}r^2 \frac{3}{5} \cdot \frac{4}{5} + \frac{5}{4}r^2 \sin^{-1}\left(\frac{3}{5}\right)$ $= \frac{3}{5}r^2 + \frac{5}{4}r^2 \sin^{-1}\left(\frac{3}{5}\right)$ $\int_0^{\frac{3r}{5}} 2r \, dx = \left[2rx \right]_0^{\frac{3r}{5}} = \frac{6r^2}{5}$ $\text{Area} = \frac{3}{5}r^2 + \frac{5}{4}r^2 \sin^{-1}\left(\frac{3}{5}\right) - \frac{6}{5}r^2$ $\text{Area} = \frac{5}{4}r^2 \sin^{-1}\left(\frac{3}{5}\right) - \frac{3}{5}r^2$ $= 0.2044r^2$		
--	--	--	--

THREE (a)(i)	Since $y = k + \frac{1}{m} \ln\left(\frac{k}{x}\right)$ $\frac{k}{x} = e^{m(y-k)}, x = ke^{-m(y-k)}$ So the area A is given by $A = \int_0^a ke^{m(k-y)} dy = ke^{mk} \int_0^a e^{-my} dy$ $= ke^{mk} \left[-\frac{1}{m} e^{-my} \right]_0^a = -\frac{k}{m} e^{mk} (e^{-ma} - 1)$ When $-\frac{k}{m} e^{mk} (e^{-ma} - 1) = \frac{p}{100} \cdot \frac{k}{m} e^{mk}$ $-(e^{-ma} - 1) = \frac{p}{100} \quad \left(\frac{k}{m} e^{mk} \neq 0 \right)$ $e^{ma} = \frac{100}{100 - p}$ $a = \frac{1}{m} \ln\left(\frac{100}{100 - p}\right).$	6	
		4	Or equivalent.

THREE (b)	$y = x \sin nx + \frac{1}{n} \cos nx$ $\frac{dy}{dx} = \sin nx + nx \cos nx - \sin nx$ $= nx \cos nx$ $I_n - I_{n-1}$ $= \int_0^{\pi} \left(\frac{1}{2}\pi - x \right) \sin \left(n + \frac{1}{2} \right) x \operatorname{cosec} \frac{1}{2} x - \left(\frac{1}{2}\pi - x \right) \sin \left(n - \frac{1}{2} \right) x \operatorname{cosec} \frac{1}{2} x \, dx$ $= \int_0^{\pi} \left(\frac{1}{2}\pi - x \right) \operatorname{cosec} \frac{1}{2} x \left(\sin \left(n + \frac{1}{2} \right) x - \sin \left(n - \frac{1}{2} \right) x \right) dx$ $= \int_0^{\pi} \left(\frac{1}{2}\pi - x \right) \operatorname{cosec} \frac{1}{2} x \left(2 \cos nx \sin \frac{1}{2} x \right) dx$ $= \int_0^{\pi} \left(\frac{1}{2}\pi - x \right) 2 \cos nx \, dx$ $= \int_0^{\pi} \pi \cos nx - 2x \cos nx \, dx$ $= \left[\frac{\pi}{n} \sin nx - \frac{2}{n^2} (nx \sin nx + \cos nx) \right]_0^{\pi} \quad \text{using the first result}$ $= \left(\frac{\pi}{n} \sin n\pi - \frac{2}{n^2} (n\pi \sin n\pi + \cos n\pi) \right) - \left(\frac{\pi}{n} \sin 0 - \frac{2}{n^2} (n\pi \sin 0 + \cos 0) \right)$	8	
	$= \left(0 - \frac{2}{n^2} (0+1) \right) - \left(0 - \frac{2}{n^2} (0+1) \right) \text{ for } n \text{ even}$ $= 0 \text{ for } n \text{ even}$ $= \left(0 - \frac{2}{n^2} (0-1) \right) - \left(0 - \frac{2}{n^2} (0+1) \right) \text{ for } n \text{ odd}$ $= \frac{2}{n^2} + \frac{2}{n^2}$ $= \frac{4}{n^2} \text{ for } n \text{ an odd number.}$ $I_0 = 0 \text{ (given)}$ $n = 1 (n \text{ odd}), \quad I_1 - I_0 = \frac{4}{1^2} \quad \text{so} \quad I_1 = \frac{4}{1^2}$ $n = 2 (n \text{ even}), \quad I_2 - I_1 = 0 \quad \text{so} \quad I_2 = I_1 = \frac{4}{1^2}$ $n = 3 (n \text{ odd}), \quad I_3 - I_2 = \frac{4}{3^2} \quad \text{so} \quad I_3 = \frac{4}{3^2} + \frac{4}{1^2}$	6	

Three (b) contd	$n = 4 \text{ (} n \text{ even)}, \quad I_4 - I_3 = 0 \quad \text{so} \quad I_4 = I_3 = \frac{4}{3^2} + \frac{4}{1^2}$ $n \text{ even, } I_n = \frac{4}{1^2} + \frac{4}{3^2} + \dots + \frac{4}{(n-1)^2} = 4 \sum_{i=1}^n \frac{1}{(i-1)^2}$ $n \text{ odd, } I_n = \frac{4}{1^2} + \frac{4}{3^2} + \dots + \frac{4}{n^2} = 4 \sum_{i=1}^n \frac{1}{i^2}$		$\sum$ notation not required.
----------------------------------	---	--	----------------------------------

and
$$\frac{dx}{dt} \cdot \frac{d^2y}{dt^2} = \frac{dy}{dt} \cdot \frac{d^2x}{dt^2}$$

Method 3

$$\begin{aligned} \frac{d^2y}{dx^2} &= \frac{d}{dx} \left(\frac{dy}{dx} \right) \\ &= \frac{d}{dx} \left(\frac{dy}{dt} \cdot \frac{dt}{dx} \right) \\ &= \frac{d}{dx} \left(\frac{dy}{dt} \right) \frac{dt}{dx} + \frac{d}{dx} \left(\frac{dt}{dx} \right) \frac{dy}{dt} \\ &= \frac{d}{dt} \left(\frac{dy}{dt} \right) \left(\frac{dt}{dx} \right)^2 + \frac{d^2t}{dx^2} \cdot \frac{dy}{dt} \\ &= \frac{d^2y}{dt^2} \left(\frac{dt}{dx} \right)^2 + \frac{d^2t}{dx^2} \cdot \frac{dy}{dt} = 0 \end{aligned}$$

but
$$\frac{dt}{dx} = \left(\frac{dx}{dt} \right)^{-1}$$

$$\begin{aligned} \frac{d^2t}{dx^2} &= - \left(\frac{dx}{dt} \right)^{-2} \frac{d^2x}{dt^2} \cdot \frac{dt}{dx} \\ &= - \frac{d^2x}{dt^2} \left(\frac{dt}{dx} \right)^3 \end{aligned}$$

hence
$$\frac{d^2y}{dt^2} \left(\frac{dt}{dx} \right)^2 = - \left(- \frac{d^2x}{dt^2} \left(\frac{dt}{dx} \right)^3 \right) \frac{dy}{dt}$$

$$\frac{d^2y}{dt^2} \cdot \frac{dx}{dt} = \frac{d^2x}{dt^2} \cdot \frac{dy}{dt}$$

Method 4

$$\frac{d^2y}{dx^2} = 0$$
 so integrating wrt x

$$\frac{dy}{dx} = k = \frac{dy}{dt} \cdot \frac{dt}{dx} \text{ and so } \frac{dy}{dt} = k \frac{dx}{dt}$$

Differentiating this wrt t

$$\frac{d^2y}{dt^2} = k \frac{d^2x}{dt^2} \text{ and } \frac{d^2y}{dt^2} = \frac{dy}{dt} \cdot \frac{dt}{dx} \cdot \frac{d^2x}{dt^2} \text{ so}$$

$$\frac{d^2y}{dt^2} \cdot \frac{dx}{dt} = \frac{d^2x}{dt^2} \cdot \frac{dy}{dt}$$

FOUR (a)(ii)	$x = a \cos t + \frac{1}{2} b \cos 2t$ $\frac{dx}{dt} = -a \sin t - b \sin 2t$ $\frac{d^2x}{dt^2} = -a \cos t - 2b \cos 2t$ $y = a \sin t + \frac{1}{2} b \sin 2t$ $\frac{dy}{dt} = a \cos t + b \cos 2t$ $\frac{d^2y}{dt^2} = -a \sin t - 2b \sin 2t$ For points of inflection $\frac{d^2y}{dx^2} = 0$, and using the result from 4(a)(i) $(-a \sin t - b \sin 2t)(-a \sin t - 2b \sin 2t)$ $-(a \cos t + b \cos 2t)(-a \cos t - 2b \cos 2t) = 0$ $a^2 \sin^2 t + 2ab \sin t \sin 2t + ab \sin t \sin 2t + 2b^2 \sin^2 2t +$ $a^2 \cos^2 t + 2ab \cos t \cos 2t + ab \cos t \cos 2t + 2b^2 \cos^2 2t = 0$ $a^2 (\sin^2 t + \cos^2 t) + 3ab (\cos t \cos 2t + \sin t \sin 2t) +$ $2b^2 (\sin^2 2t + \cos^2 2t) = 0$ $a^2 + 3ab \cos t + 2b^2 = 0$ $\cos t = \frac{-(a^2 + 2b^2)}{3ab}$	6	
FOUR (b)	$\frac{dx}{dy} \cdot \frac{d^2y}{dx^2} = k \frac{dy}{dx}$ $\text{so } \frac{d\left(\frac{dy}{dx}\right)}{dx} = k \left(\frac{dy}{dx}\right)^2 \quad \text{Let } z = \frac{dy}{dx}$ $\frac{dz}{dx} = kz^2$ $\int \frac{1}{z^2} dz = k \int dx$ $-\frac{1}{z} = kx + C \quad \text{but } z = 1 \text{ when } x = 0 \text{ (given), so } C = -1$	8	

FOUR (b) contd	$-\frac{1}{z} = kx - 1$ $\frac{dy}{dx} = \frac{1}{1 - kx}$ $y = \int \frac{1}{1 - kx} dx = -\frac{1}{k} \ln 1 - kx + C \quad \text{but when } x = 0, y = 1 \text{ (given)}$ so $C = 1$ $y = 1 - \frac{1}{k} \ln 1 - kx $ and when $y = 2$, $1 = -\frac{1}{k} \ln 1 - kx$, $1 - kx = e^{-k}$, $x = \frac{1}{k}(1 - e^{-k})$ so when $y = 2$, $\frac{dy}{dx} = \frac{1}{1 - kx} = \frac{1}{e^{-k}} = e^k$. Or Let $\frac{dy}{dx} = p$, then $\frac{d^2y}{dx^2} = \frac{dp}{dy} p$ and so for $\frac{dx}{dy} \cdot \frac{d^2y}{dx^2} = k \frac{dy}{dx}$ $p \frac{dp}{dy} = kp^2$ $\int \frac{1}{p} dp = k \int dy \quad \text{and}$ $\ln p = ky + C$ $p = Ae^{ky}$ but when $y = 1$, $p = \frac{dy}{dx} = 1$, so $A = e^{-k}$ $p = e^{k(y-1)} \quad \text{and so when } y = 2 \quad p = \frac{dy}{dx} = e^k$	6	
---	---	-----------------	--

SIX (a) Contd	$\frac{2(x-2h)}{a^2} - \frac{2(y-h)}{4a^2} \frac{dy}{dx} = 0$ $\frac{dy}{dx} = \frac{4(x-2h)}{y-h}$ and when $x = p, y = q$ $\frac{dy}{dx} = \frac{4(p-2h)}{q-h} \text{ and so}$ $\frac{4(p-2h)}{q-h} = 4$ $p-2h = q-h, \quad p-q = h.$	2 2	Or differentiating the original form. Gradient must be correct.
SIX (b)	$G = (-ae, 0) = (-a\sqrt{5}, 0)$; $F = (ae, 0) = (a\sqrt{5}, 0)$ for a standard hyperbola, so Either: We can work with these points and then translate the line by $\begin{pmatrix} 2h \\ h \end{pmatrix}$ For A: $\frac{(a\sqrt{5})^2}{a^2} \cdot \frac{y^2}{4a^2} = 1, \quad \frac{y^2}{4a^2} = 5 - 1 = 4, \quad y^2 = 16a^2, \quad y = \pm 4a$ So $A = (a\sqrt{5}, 4a)$ and the line AG is: $y - 0 = \frac{4a}{a\sqrt{5} + a\sqrt{5}}(x + a\sqrt{5})$ $y = \frac{2}{\sqrt{5}}(x + a\sqrt{5})$ then translating the line by $\begin{pmatrix} 2h \\ h \end{pmatrix}$ $y - h = \frac{2}{\sqrt{5}}(x - 2h + a\sqrt{5}) \quad \text{or}$ $\sqrt{5}y = 2x + (\sqrt{5} - 4)h + 2a\sqrt{5}.$ Or: with a translation $\begin{pmatrix} 2h \\ h \end{pmatrix}$ of the points first we get $G = (2h - a\sqrt{5}, h), \quad F = (2h + a\sqrt{5}, h)$ Hence for A $\frac{(2h + a\sqrt{5} - 2h)^2}{a^2} - \frac{(y - h)^2}{4a^2} = 1$	8 4	Or equivalent.

SIX (c) contd	$(mx - y) = \frac{b^2}{y_1} = \frac{b^2}{\pm \frac{b^2}{\sqrt{m^2 a^2 - b^2}}} = \pm \sqrt{m^2 a^2 - b^2}$ and $y = mx \pm \sqrt{m^2 a^2 - b^2}$.		
SIX (d)	This question is equivalent to the problem obtained by translating the hyperbola and using $\frac{x_1^2}{a^2} - \frac{y_1^2}{4a^2} = 1 \quad \text{and a tangent through the point } \left(\frac{2a}{3}, 0\right).$ Using the answer to part (c) and the point $\left(\frac{2a}{3}, 0\right)$ $0 = \frac{2am}{3} \pm \sqrt{m^2 a^2 - 4a^2}$ $\frac{4a^2 m^2}{9} = (m^2 - 4)a^2$ $5m^2 = 36, \quad m = \pm \frac{6}{\sqrt{5}} \quad \text{and we want the + sign as asked so}$ $m = \frac{6}{\sqrt{5}}.$ The asymptote is $y = \frac{bx}{a} = \frac{2ax}{a} = 2x$, with gradient 2. So the angle α between the two lines is given by $\tan \alpha = \frac{m_1 - m_2}{1 + m_1 m_2} = \frac{\frac{6}{\sqrt{5}} - 2}{1 + 2 \cdot \frac{6}{\sqrt{5}}} = \frac{6 - 2\sqrt{5}}{\sqrt{5} + 12}$ $= \frac{(6 - 2\sqrt{5})(\sqrt{5} - 12)}{(\sqrt{5} + 12)(\sqrt{5} - 12)} = \frac{30\sqrt{5} - 82}{-139} = \frac{82 - 30\sqrt{5}}{139}$ and $\alpha = \tan^{-1} \left(\frac{2(41 - 15\sqrt{5})}{139} \right).$	8 6	

An aggregate mark of 120 from six questions was used in Scholarship Calculus. In 2005, candidates who achieved 47–120 were awarded outstanding scholarship, and candidates who achieved 20–46 were awarded scholarship.