



For Supervisor's use only

3

90522



NEW ZEALAND QUALIFICATIONS AUTHORITY
MANA TOHU MĀTAURANGA O AOTEAROA



National Certificate of Educational Achievement
TAUMATA MĀTAURANGA Ā-MOTU KUA TAEA

Level 3 Physics, 2006

90522 Demonstrate understanding of atoms, photons and nuclei

Credits: Three

9.30 am Monday 20 November 2006

Check that the National Student Number (NSN) on your admission slip is the same as the number at the top of this page.

You should answer ALL the questions in this booklet.

For all numerical answers, full working must be shown, and the answer must be rounded to the correct number of significant figures and given with an SI unit.

For all 'describe' or 'explain' questions, the answers should be written or drawn clearly with all logic fully explained.

Formulae you may find useful are given on page 2.

If you need more space for any answer, use the page(s) provided at the back of this booklet and clearly number the question.

Check that this booklet has pages 2–8 in the correct order and that none of these pages is blank.

YOU MUST HAND THIS BOOKLET TO THE SUPERVISOR AT THE END OF THE EXAMINATION.

For Assessor's use only		Achievement Criteria	
Achievement		Achievement with Merit	Achievement with Excellence
Identify or describe aspects of phenomena, concepts or principles.	<input type="checkbox"/>	Give explanations in terms of phenomena, concepts, principles and/or relationships.	<input type="checkbox"/>
Solve straightforward problems.	<input type="checkbox"/>	Solve problems.	<input type="checkbox"/>
Overall Level of Performance (all criteria within a column are met)			<input type="checkbox"/>

You are advised to spend 35 minutes answering the questions in this booklet.

Assessor's
use only

You may find the following formulae useful.

$$E = hf$$

$$hf = \phi + E_K$$

$$E = \Delta mc^2$$

$$E_n = -\frac{hcR}{n^2}$$

$$\frac{1}{\lambda} = R\left(\frac{1}{S^2} - \frac{1}{L^2}\right)$$

$$E_p = qV$$

$$v = f\lambda$$

QUESTION ONE: NUCLEAR REACTIONS

Mass of nuclei:

neutron: $1.67492 \times 10^{-27} \text{ kg}$

proton: $1.67353 \times 10^{-27} \text{ kg}$

deuterium: $3.34449 \times 10^{-27} \text{ kg}$

tritium: $5.00827 \times 10^{-27} \text{ kg}$

helium-4: $6.64648 \times 10^{-27} \text{ kg}$

lithium-6: $9.98835 \times 10^{-27} \text{ kg}$

Speed of light = $3.00 \times 10^8 \text{ m s}^{-1}$



Three bottles of water and some rocks can provide, in theory, enough energy for a family for one year. The water and rocks can be used to obtain the raw materials for a thermonuclear reaction that can take place between deuterium and tritium.

Tritium can be made from lithium ${}^6_3\text{Li}$, which can be extracted from the rocks.

- (a) Show that the mass deficit of a lithium nucleus is $5.700 \times 10^{-29} \text{ kg}$.


- (b) Calculate the binding energy per nucleon for the lithium nucleus.

binding energy per nucleon = _____

- (c) State how the binding energy per nucleon can indicate the **stability** of a nucleus.

Assessor's
use only

Deuterium (hydrogen-2) can be extracted from the water. Thermonuclear reactors heat a mixture of deuterium and tritium to 100 million degrees Celsius to produce the reaction illustrated below.



The nuclear equation for this reaction is: ${}^2_1\text{H} + {}^3_1\text{H} \rightarrow {}^4_2\text{He} + {}^1_0\text{n}$

- (d) Calculate the amount of energy produced in this reaction.

energy = _____

- (e) Explain why it is necessary for the temperature to be **so high** for this reaction to occur.

QUESTION TWO: SOLAR POWERAssessor's
use onlyRydberg's constant = $1.097 \times 10^7 \text{ m}^{-1}$ Planck's constant = $6.63 \times 10^{-34} \text{ J s}$ Speed of light = $3.00 \times 10^8 \text{ m s}^{-1}$

Nuclear reactions in the Sun produce light. The main element in the Sun is hydrogen. The spectrum of hydrogen can be observed in the laboratory with a hydrogen discharge tube.

The visible lines in the hydrogen spectrum are called the Balmer series and are described by the formula:

$$\frac{1}{\lambda} = R \left(\frac{1}{S^2} - \frac{1}{L^2} \right)$$

where $S = 2$.

- (a) Calculate the wavelength of the lowest frequency line in the Balmer series ($L = 3$). Give the answer to the correct number of significant figures.

wavelength =

- (b) Explain how light of **this particular** frequency is produced in the hydrogen atom.

- (c) An electron in the 6th excited state ($L = 7$) returns to the ground state in two jumps. It releases one photon with a wavelength of 2.165×10^{-6} m.

What is the wavelength of the second photon?

wavelength = _____


- (d) The Sun emits **all** wavelengths. However, when a solar spectrum is observed on Earth, it contains black lines that correspond to missing wavelengths.

Give an explanation, in terms of energy absorption by electrons, for why some of the wavelengths of light in the solar spectrum are missing when the light reaches Earth.

QUESTION THREE: NIGHT VISION CAMERAAssessor's
use only

Planck's constant = $6.63 \times 10^{-34} \text{ J s}$

A night vision camera, like the one shown below, detects low levels of light on the photo-cathode, which releases a few electrons. A photomultiplier increases the number of electrons, which then hit the screen to produce an image.



Source <http://www.lancs.ac.uk/ug/jackson2/>

- (a) Name the **effect** that causes electrons to be released by the photo-cathode.
- _____
- (b) The photo-cathode material of this night vision camera prevents it detecting infrared radiation. State **why** this is so.
- _____
- _____

The photo-cathode is made of a material that has a work function of $2.58 \times 10^{-19} \text{ J}$.

- (c) Calculate the lowest frequency of light that could release a photoelectron.
- _____
- _____
- _____

frequency = _____

(e) Calculate the threshold frequency for the material.

threshold frequency = _____

**Extra paper for continuation of answers if required.
Clearly number the question.**

Assessor's
use only

Question
number